LOCAL NAMES

Arabic (arrad); Swahili (mgunga); Tigrigna (seraw)

BOTANIC DESCRIPTION

Acacia etbaica is a tree or shrub (min. 2) 2.5-12 m tall; trunk distinct with flattened or round crown in older trees; bark pale brown, brownish-black, reddish-grey or dark grey, smooth or fissured; spines in pairs at the nodes, brownish-white, of 2 kinds-small and recurved, to 7 mm long and long to 6 mm; slash fibrous and creamy white; twigs red-brown.

Leaves have 1-9 pairs of pinnae; leaflets in 4-35 pairs, pubescent, 0.5-4 x 0.2-1.3 mm; petiole 0.3-1 cm; petiole and rachis glands absent.

Inflorescences capitate, on axillary peduncles 0.7-4 cm long; involucel 1/3 to 2/3 up peduncle, or sometimes near the base; flowers white or cream; calyx 0.4-1 mm long, glabrous apices of lobes puberulous; corolla 2-3 mm long.

Fruit a pod, linear-oblong, purple-brown to red brown, shiny, straight, 2-12 \times 0.6-2.2 cm, often attenuated at the base, glabrous or puberulous, with fine oblique or longitudinal veins, dehiscent; seeds about 8, brown or olive-brown, elliptic, flattened, 5.5 \times 3.5-8 mm.

A. etbaica has a number of subspecies namely: subsp. etbaica, mainly found in Sudan and Somalia, subsp. uncinata Brenan found in Somalia, Uganda and Kenya and subsp. platycarpa Brenan found in Kenya and Tanzania. A. etbaica is also closely related to A. reficiens ssp. misera and A. elatior.

The generic name 'acacia' comes from the Greek word 'akis', meaning a point or a barb.

BIOLOGY

A. etbaica is hermaphroditic.

ECOLOGY

A. etbaica occurs in dry bushland, thickets, semi-desert scrub and wooded grasslands.


BIOPHYSICAL LIMITS

Altitude: 0-1 800 m, Mean annual rainfall: 200 -1400mm, Mean annual temperature: 22.3 deg.C

DOCUMENTED SPECIES DISTRIBUTION

Native: Eritrea, Ethiopia, Kenya, Somalia, Sudan, Tanzania, Uganda

Exotic:



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

Fabaceae - Mimosoideae

PRODUCTS

Fuel: A. etbaica is a source of good firewood.

Timber: Provides the pillars and beams to hold the heavy earthen roofs of houses in northern Ethiopia.

Medicine: The bark is chewed as a stimulant and is also used in the treatment of gonorrhoea

Fabaceae - Mimosoideae

GERMPLASM MANAGEMENT Seed storage behaviour is orthodox.

Fabaceae - Mimosoideae

FURTHER READNG

Beentje HJ. 1994. Kenya trees, shrubs and lianas. National Museums of Kenya.

Bein E. 1996. Useful trees and shrubs in Eritrea. Regional Soil Conservation Unit (RSCU), Nairobi, Kenya.

Coe M and Beentje H. 1991. A field guide to the Acacias of Kenya. Oxford University Press.

Dale IR, Greenway PJ. 1961. Kenya trees and shrubs. Buchanan's Kenya Estates Ltd.

Eggeling. 1940. Indigenous trees of Uganda. Govt. of Uganda.

El Amin HM. 1973. Sudan acacias. Forest Research Institute Publishing Section Information Department.

Hedberg I and Edwards S (eds.). 1989. Flora of Ethiopia Volume 3: Pittosporaceae to Araliaceae. The National Herbarium, Biology Department, Addis Ababa University and The Department of Systematic Botany, Uppsala University, Sweden.

Hong TD, Linington S, Ellis RH. 1996. Seed storage behaviour: a compendium. Handbooks for Genebanks: No. 4. IPGRI.

Kokwaro JO. 1976. Medicinal plants of East Africa. East African Literature Bureau.

SUGGESTED CITATION

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 Agroforestree Database:a tree reference and selection guide version 4.0 (http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp)