

Pometia pinnata

kasai

J.R. Forster & J.G. Forster

Sapindaceae

LOCAL NAMES

Burmese (paga-nyet-su ava); Filipino (tugau); Indonesian (leungsir, ihi mendek, tawan, kayu sapi); Lao (Sino-Tibetan) (kwaang, chieng dong); Malay (kasai daun besar, langsir, sibu); Thai (saen ta lom, sai); Trade name (kasai); Vietnamese (tr[uw][owf]ng m[aa]t, m[aws]c ken)

BOTANIC DESCRIPTION

Pometia pinnata is a medium-sized to fairly large tree up to 40(-47) m tall, with bole up to 100(-140) cm in diameter.

Leaves with 3-13 pairs of leaflets having dentate margins, and alternate secondary veins ending in a tooth.

Inflorescence and calyx often hairy.


P. pinnata is an extremely variable species. No less than 8 forms have been recognized and given taxonomic status, but because still other and intermediate forms are encountered, this distinction is probably only useful in e.g. breeding. For reasons of completeness the 8 forms distinguished are given here : f. *pinnata* (synonym: *P. coriacea* Radlk., 1913); f. *acuminata* (Hook.f.) Jacobs (synonyms: *P. acuminata* (Hook.f.) Radlk., 1877, *P. annamica* Gagn., 1947); f. *alnifolia* (Blume) Jacobs (synonym: *P. alnifolia* (Blume) Radlk., 1877); f. *cuspidata* (Blume) Jacobs; f. *glabra* (Blume) Jacobs (synonym: *P. pinnata* var. *javanica* Koord. & Valetton, 1903); f. *macrocarpa* (Kurz) Jacobs (synonym: *P. macrocarpa* Kurz, 1875); f. *repandra* Jacobs; f. *tomentosa* (Blume) Jacobs (synonym: *P. tomentosa* (Blume) Teijsm. & Binnend., 1866).

BIOLOGY

Bisexual and male flowers are reported to occur on a single tree of *P. pinnata*, but the structurally hermaphrodite flowers are functionally female with no anther dehiscence. Both cross-fertilization and self-fertilization occur. Usually there are 3-4 times as many male as female flowers. Dispersal of the fruits is probably mostly by bats and birds.



Ton fruit (French B.)



Ton leaf (French B.)

kasai

ECOLOGY

P. pinnata occurs in primary and secondary forests.

BIOPHYSICAL LIMITS

Mean annual rainfall: 1 000-4 600 mm, Mean annual temperature: 20-29 deg.C

DOCUMENTED SPECIES DISTRIBUTION

Native: India, Papua New Guinea, Philippines, Solomon Islands, Sri Lanka

Exotic:



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

kasai

PRODUCTS

Food: The tree is sometimes cultivated for its fruits which can have a thick and sweet-tasting arilode. The seeds are also edible.

Fuel: The wood is also used for charcoal production.

Timber: Heartwood light to dark red, medium dark red-brown, sometimes purplish, sometimes with dark coloured ribbons on radial and tangential sections, not always well demarcated from the pink or buff-coloured sapwood. Grain straight or slightly interlocked. Texture rather coarse but even; wood surface glossy. The wood density *P. pinnata* is 500-990 kg/m cubic at 15% moisture content. The timber is used as kasai.

Medicine: The leaves and bark are used medicinally.

SERVICES

Pometia pinnata

kasai

J.R. Forster & J.G. Forster

Sapindaceae

TREE MANAGEMENT

GERMPLASM MANAGEMENT

PESTS AND DISEASES

FURTHER READNG

- Allen BJ. 1985. Dynamics of fallow successions and introduction of robusta coffee in shifting cultivation areas in the lowlands of Papua New Guinea. *Agroforestry systems*. 3(3): 227-238.
- Backer CA, Bakhuizen van den Brink RC. 1963-1968. *Flora of Java*. 3 volumes. Noordhoff, Groningen, the Netherlands. pp. 20-35.
- Brown WH. 1978. *Timbers of the world*. Vol 3. Southern Asia, 99 pp.; vol. 4, South East Asia, 82 pp.; vol. 8, Australasia, 93 pp. Timber Research and Development Association (TRADA), High Wycombe.
- Browne FG. 1955. *Forest trees of Sarawaka and Brunei and their products*. Government Printing Office, Kuching. 369 pp.
- Budgen B. 1981. The shrinkage and density of some Australian and South-east Asian Timbers. Technical Paper (2nd series) No 38. Division of Building Research, CSIRO, Melbourne. 38 pp.
- Burkill IH. 1966. *A dictionary of the economic products of the Malay Peninsula*. Revised reprint. 2 volumes. Ministry of Agriculture and Co-operatives, Kuala Lumpur, Malaysia. Vol. 1 (A-H) pp. 1-1240. Vol. 2 (I-Z) pp. 1241-2444.
- Burslem DFRP & Whitmore TC. 1996. *Silvics and wood properties of the common timber tree species on Kolombangara*. Oxford Forestry Institute Tropical Forestry Paper No. 34.
- Chudnoff M. 1984. *Tropical timbers of the world*. Agricultural Handbook 607. USDA Forest Service, Washington, DC. 464 pp.
- Corner EJH. 1988. *Wayside trees of Malaya*, 3rd Edition. 2 volumes. The Malayan Nature Society, Kuala Lumpur, Malaysia. 744p.
- de Guzman ERM, Umali & Sotalbo ED. 1986. *Guide to Philippine flora and fauna*. Vol. 3: dipterocarps, non-dipterocarps. Natural Resources Management Centre, Ministry of Natural Resources & University of the Philippines, Manila 414 pp.
- Desch HE. 1954. *Manual of Malayan timbers*. Malayan Forest Records No 15. 2 volumes. Malaya Publishing House Ltd., Singapore. 726 pp.
- Eddowes PJ. 1977. *Commercial timbers of Papua New Guinea, their properties and uses*. Forest Products research Centre, Department of Primary Industry, Port Moresby. 195 pp.
- Elmqvist T, Rainey WE, Pierson ED and Cox PA. 1994. Effects of tropical cyclones Ofa and Val on the structure of Samoan lowland rain forest. *Biotropica*. 26:384-391.
- Forest Industries Council of Papua New Guinea. 1982. *Papua New Guinea timbers*. Technical data. Port Moresby. 35 pp
- Forestry Division, Ministry of Natural Resources. 1979. *major species*. Timber booklet No. 1. Solomon Island Timbers. Government Printing Office, Honiara. 84 pp.
- Frenaman KW. 1967. *Data on certain B.S.I.P. timber species*. Technical Note for department of the British Solomon Islands Protection No 9/67.
- Fundter JM, de Graaf NR, & Hilderbrand JW. 1989. *Pometia pinnata* J.R. & G. Forst. In: Westphal, E. & Jensen, PCM. (Editors): *Plant resources of South-East Asia*. A selection. Pudoc, Wageningen. Pp. 230-232.
- Ginoga B, Hadjib N and Karnasudirdja S. 1982. *Sifat Fisis dan Mekanis Beberapa Jenis Kayu Indonesia Bagian 10*. Laporan BPHH No. 162. Bogor p:7-21
- Gosh SS, Ramesh Rao K & Purkayastha. (Eds). 1963. *Indian woods, their identification, properties and uses*. Vol. 2, Linaceae to Moringaceae. Manager of Publications, Delhi. 386 pp.
- Grewal GS. 1979. *Air-seasoning properties of some Malaysian timbers*. Malaysian Forest Service Trade Leaflet No 41. Malaysian timber Industry Board, Kuala Lumpur. 26 pp.
- Ha CO, et al. 1988. Reproductive patterns of selected understorey trees in the Malaysian rain forest: the sexual species. *Botanical Journal of the Linnaean Society*. 97: 295-316.
- Henderson CP, Hancock IR. 1989. *A guide to the useful plants of the Solomon Islands*. Research Department, Ministry of Agriculture and Lands, Honiara. 82 pp.
- Jackson WF. 1965. *The durability of Malaysian timbers*. Malaysian Forest Service Trade Leaflet No 28. Malaysian Timber Industry Board, Kuala Lumpur. 11 pp.
- Kloot NH & Bolza E. 1977. *Properties of timbers imported into Australia*. Technical Paper No 17 (2nd Series). Division of Building Research, CSIRO, Melbourne. 81 pp.

Lauricio FM & Bellosilo SB. 1963/64. The mechanical and related properties of Philippine woods. Philippine Limberman. 10: 49-56.

Lemmens RHMJ, Soerianegara I, Wong WC (eds.). 1995. Plant Resources of South-east Asia. No 5(2). Timber trees: minor commercial timbers. Backhuys Publishers, Leiden.

Lopez DT. 1978. The resistance to splitting in nailing some Malaysian timbers. Malayan Forestry Service Trade Leaflet No 39. Malaysian Timber Industry Board, Kuala Lumpur. 12 pp.

Martawijaya A & Barly. 1982. Resistance of Indonesian timbers to impregnation with CCA preservative. Pengumuman No 5. Lembaga Penelitian Hasil Hutan, Bogor. 18 pp.

Martin FW, Campbell CW & Ruberte RM. 1987. Perennial edible fruits of tropics: an inventory. US Department of Agriculture, Agriculture Handbook No. 642. 252 pp.

Ngok WC. 1985. Malaysian timbers for plywood manufacture. Malaysian Forest Service Trade Leaflet No 94. Malaysian timber Industry Board, Kuala Lumpur. 14 pp.

Nguyen Ngoc Chinh, et al. 1996. Vietnam forest trees. Hanoi, Vietnam: Agricultural Publishing House. 788p.

Seeber G, Weidelt HJ, Banaag VS. 1985. Dendrological characters of important forest trees from eastern Mindanao. 2nd edition. Eschborn, Germany.

Soerianegara I, Lemmens RHMJ (eds.). 1993. Plant Resources of South-East Asia. No. 5(1): Timber trees: major commercial timbers. Backhuys Publishers, Leiden.

Uriarte MT. 1994. Air-pollution resistant species recommended for urban areas (A. Highly resistant species). Techno-Info Series, 4(1) Jan-March. Quezon City, Philippines: ERDS-NCR-DENR.

Walter A, Sam C. 2002. Fruits of Oceania. ACIAR Monograph No. 85. Canberra. 329 pp.

Whistler. 1992. Vegetation of Samoa and Tonga. Pacific Science. 46:159-178.

Whitmore TC, Tantra IGM. 1986. Tree flora of Indonesia. Checklist for Sumatra. Bogor, Indonesia: Forest Research and Development Centre.

Williams MD, Logan AF, et al. (1984). The utilization of defective *Pometia pinnata* for pulpwood. *Appita*. 37(7): 587-591.

Wong TM. 1982. A dictionary of Malayan Timbers. Malayan Forest Records No 30. Forest Department, Kuala Lumpur. 529 pp.

SUGGESTED CITATION

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 Agroforestry Database: a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)