

LOCAL NAMES

Afrikaans (rivierboontjie); Amharic (girangire); Arabic (sesaban); Bengali (jainti,jayant); Burmese (yay-tha-kyee,yethugyi); English (common sesban,Egyptian rattle pod,frother,sesbania,sesban,river bean); Filipino (katuray,katodai); Hindi (jait,jainti,rawasan); Indonesian (janti,puri,jayanti); Javanese (janti); Khmer (snaô kôôk); Lao (Sino-Tibetan) (sapao lom); Luganda (muzimbandeya,mubimba); Sanskrit (jayantika,jayanti); Spanish (Añil francés,tamarindillo); Tamil (karunchembai,chithagathi,champai); Thai (sami,saphaolom); Vietnamese (dien-dien); Zulu (umQambuqweqwe,umsokosoko)

BOTANIC DESCRIPTION

Sesbania sesban is a narrow-crowned, deep-rooting single or multi stemmed shrub or small tree, 1-7 m tall. The trees usually have a main stem but may develop many side branches if widely spaced. The many branches give the tree a shrubby appearance, often tending towards a spreading habit due to its wide branching angle (45-60 deg. Mostly).

Leaves paripinnate, long, narrow; leaflets in many pairs, rounded or oblong, usually asymmetric at the base, often glaucous; stipules minute or absent.

Flowers attractive, yellow, red, purplish, variegated or streaked, seldom white, large or small on slender pedicels, solitary or paired in short axillary racemes, usually unpleasantly scented; all petals long clawed, standard orbicular or obovate.

Pods pale yellow, linear, usually 10-20 cm long, cylindrical or compressed, rarely oblong; up to 40 seeds are found in a pod; seeds oblong or subquadrate, brown or dark green mottled with black.

Two subspecies are recognized within *S. sesban*, namely ssp. *punctata* (restricted to northern portions of sub-Saharan Africa) and ssp. *sesban*.

BIOLOGY

S. sesban is assumed to be largely out-crossing, however interspecific hybridization is reported with *S. goetzei*; the carpenter bee is its main pollinator. Flowering starts shortly after the onset of the rains (in areas where there are 2 rainy seasons, it flowers and sets fruit twice). Pods are indehiscent and do not shed their seeds until well after pod maturity.



S. sesban flowers (Anthony Simons)



S. sesban Mesoplatys beetle attack. (Anthony Simons)



S. sesban pods. (Anthony Simons)

ECOLOGY

S. sesban grows well in the subtropics and is significant in extending the nitrogen-fixing forage trees into cooler, higher elevation regions of the tropics. It has outstanding ability to withstand waterlogging and is ideally suited to seasonally flooded environments. When flooded, it initiates floating, adventitious roots and protects its stems, roots and nodules with spongy, aerenchyma tissue. It is common along streams, swamp banks and moist and inundated bottomlands. *S. sesban* shows some tolerance to moisture stress and tolerates soil alkalinity and salinity to a considerable degree.

BIOPHYSICAL LIMITS


Altitude: 100-2300 m, Mean annual temperature: (10 min.) 18-23 (45 max.) deg. C, Mean annual rainfall: 500-2000 mm

Soil type: Tolerates seasonal or permanently waterlogged soils as well as saline, acidic and alkaline soils.

DOCUMENTED SPECIES DISTRIBUTION

Native: Chad, Egypt, Kenya, Uganda

Exotic: Angola, Australia, Benin, Botswana, Burkina Faso, Burundi, Cameroon, Cape Verde, Central African Republic, China, Congo, Cook Islands, Cote d'Ivoire, Democratic Republic of Congo, Djibouti, Equatorial Guinea, Eritrea, Ethiopia, Fiji, French Polynesia, Gabon, Gambia, Ghana, Guinea, Guinea-Bissau, India, Indonesia, Iraq, Kiribati, Laos, Lesotho, Liberia, Madagascar, Malawi, Mali, Marshall Islands, Mauritania, Mozambique, Myanmar, Namibia, New Caledonia, Niger, Nigeria, Norfolk Island, Pakistan, Papua New Guinea, Philippines, Rwanda, Samoa, Sao Tome et Principe, Seychelles, Sierra Leone, Solomon Islands, Somalia, South Africa, Sudan, Swaziland, Tanzania, Thailand, Togo, Tonga, United States of America, Vanuatu, Vietnam, Zambia, Zimbabwe



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

PRODUCTS

Food: *S. sesban* flowers are edible and are included perhaps as a decorative or festive ingredient in foods such as omelettes.

Fodder: The tree has a high percentage of foliage nitrogen and is an excellent supplement to protein-poor roughage in ruminant diets. Ruminants readily eat leaves and young branches. The crude protein content of the foliage is generally greater than 20% and often above 25%. In vitro dry-matter digestibility is 75%. Nylon-bag dry-matter digestibility of dried leaf of *S. sesban* is 90.7% and nitrogen digestibility is 96.7%. These characteristics, together with the generally low crude fibre content and high phosphorous levels, indicate the potential of the species as a high-quality forage source.

When grazed, the brittle tree may break too easily and expose the tree to fungal attack. It has been successfully fed as a sole diet to goats and as a supplement to low-quality forage for sheep.

Fuel: *S. sesban* is popular for firewood and charcoal because it produces a high woody biomass in a short time, which, although soft, is relatively smokeless, quick kindling and hot burning. The calorific yield for a 3-year-old tree is approximately 4350 kcal/kg.

Fibre: *S. sesban* is used for making ropes and fishnet and has potential for pulpwood production.

Gum or resin: *S. sesban* seeds and bark produce gum.

Poison: The saponin, stigmasta-galactopyranoside, which is isolated from the seeds, has glucuronide derivatives of oleanolic acid, which has molluscicidal activity against *Biophalaria glabrata*, one of the known snail vectors of schistosomiasis. The saponin also shows spermicidal and haemolytic activity. Using *S. sesban* leaf meal in poultry diets (as 10% of the diet) is fatal to young chicks, and the provision of either cholesterol or sitosterol with the diet significantly improves chick survival.

Medicine: Fresh *S. sesban* roots and leaves are used to treat scorpion stings, boils and abscesses. The Hausa of Ghana use decoctions of leaves as a drench for cattle to repel tsetse fly. Among the Haya people of Tanzania, it is used to treat sore throat, gonorrhoea, syphilis, spasmodic fits in children and jaundice during pregnancy. The leaves are used in some countries as a tea and are considered to have antibiotic, anthelmintic, antitumour and contraceptive properties. Oil from the seeds is accorded special properties in ayurvedic medicine and is reported to have bactericidal, cardiac depressant and hypoglycaemic actions.

SERVICES

Shade or shelter: *S. sesban* has been used to shade coffee, tea and cocoa. It has also been used as a windbreak for bananas, citrus and coffee.

Soil improver: *S. sesban* will increase soil nitrogen through symbiotic interaction with bacteria, has the ability to stabilize soil, and in Asia has been used as green manure for rice. Its branches have been used as mulch and leaves as a green manure. *S. sesban* improves soil fertility in a short-term rotation fallow and is useful in combating striga weed (*Striga hermonthica*). Some studies indicate that in 1 year a *S. sesban* fallow can increase maize yields from 2 to 4 t/ha without application of nitrogen fertilizer.

Intercropping: *S. sesban* is a promising shrub for alley cropping because it is easy to establish, it grows rapidly, coppices readily and provides mulch of high nutrient content (particularly N). In some climates, such as in the highlands of Kenya, it may have a sparse canopy, and weed competition can be a problem. This characteristic makes *S. sesban* a good intercrop.

Boundary or barrier or support: Suitable for use as live trellises for pepper.

TREE MANAGEMENT

One of the major advantages of sesbania over other forage trees and shrubs is its rapid early growth rate, which can be exploited by intercropping it with other slower establishing species for earlier yields. In India, it has been reported to attain a height of 4-5 m in 6 months. *S. sesban* thrives under repeated cuttings and coppices readily, with many branches arising from the main stem below cutting height. Cutting frequencies are generally 3-4 cuts/annum, but up to 8 cuts are made in some areas. Yields have ranged from 4 to 12 t/ha dry matter per year, depending on location. Cutting height can also influence yield, with cutting heights of 50-76 cm favouring plant survival and productivity.

The rhizobium requirements of *S. sesban* vary. There is a host-strain interaction, and different accessions of *S. sesban* require different strains of bacteria.

GERMPLASM MANAGEMENT

Seed storage behaviour is orthodox. Viability can be maintained for 2 years in open storage at room temperature. There are 85 000-100 000 seeds/kg.

PESTS AND DISEASES

S. sesban is attacked by nematodes, insects, fungi and viruses. The leaf-eating beetle *Mesoplatys ochroptera* can completely defoliate *S. sesban*, leading to mortality. Caterpillars, Hymenoptera, and stem borers are normally associated with *S. sesban*. Some potentially destructive root-knot nematodes have been recorded in India as associated with *S. sesban*.

FURTHER READING

- Albrecht J. ed. 1993. Tree seed hand book of Kenya. GTZ Forestry Seed Center Muguga, Nairobi, Kenya.
- Allen ON, Allen EK. 1981. The Leguminosae. A source book of characteristics, uses and nodulation. Macmillan, London.
- Anon. 1986. The useful plants of India. Publications & Information Directorate, CSIR, New Delhi, India.
- Beentje HJ. 1994. Kenya trees, shrubs and lianas. National Museums of Kenya.
- Bekele-Tesemma A, Birnie A, Tengnas B. 1993. Useful trees and shrubs for Ethiopia. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Chirwa TS, Mafongoya PL, Mbewe DN. 2004. Changes in soil properties and their effects on maize productivity following *Sesbania sesban* and *Cajanus cajan* improved fallow systems in eastern Zambia: Biology and Fertility of Soils. 40(1):20-27.
- Coates-Palgrave K. 1988. Trees of southern Africa. C.S. Struik Publishers Cape Town.
- Dale IR, Greenway PJ. 1961. Kenya trees and shrubs. Buchanan's Kenya Estates Ltd.
- Desaeger J, Rao MR. 2001. Effect of field establishment methods on root-knot nematode *Meloidogyne* spp. infection and growth of *Sesbania sesban* in western Kenya: Crop Protection. 20(1):31-41.
- Eggeling. 1940. Indigenous trees of Uganda. Govt. of Uganda.
- Evans DO, Macklin B. 1990. Perennial *Sesbania* species in agroforestry systems. Proceedings of a workshop, 27-31 March 1989. Nitrogen Tree Fixing Association, Paia, Hawaii, Nairobi, Kenya.
- Evans DO, Rotar P. 1987. *Sesbania* in agriculture. Westview Tropical Agricultural Series 8. Westview Press, Boulder, Colorado.
- Gutteridge RC and Shelton HM (eds.). 1994. Forage Tree Legumes in Tropical Agriculture. CAB International, Wallingford, UK.
- Hong TD, Linington S, Ellis RH. 1996. Seed storage behaviour: a compendium. Handbooks for Genebanks: No. 4. IPGRI.
- ICRAF. 1992. A selection of useful trees and shrubs for Kenya: Notes on their identification, propagation and management for use by farming and pastoral communities. ICRAF.
- Jamnadass R, Hanson J, Poole J, Hanotte O, Simons AJ, Dawson IK. 2005. High differentiation among populations of the woody legume *Sesbania sesban* in sub-Saharan Africa: implications for conservation and cultivation during germplasm introduction into agroforestry systems: Forest Ecology and Management. 210:225-238.
- Katende AB et al. 1995. Useful trees and shrubs for Uganda. Identification, Propagation and Management for Agricultural and Pastoral Communities. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Kwesiga FR. 1992. The importance of *Sesbania sesban* and close relatives in agroforestry. The ICRAF perspective. Paper presented at the planning meeting for collection of *Sesbania sesban* in Southern Africa, Lilongwe, Malawi, 23-26 Nov 1992. ICRAF.
- Leakey RRB, Newton AC. 1994. Tropical trees: the potential for domestication and rebuilding of forest resources. HMSO, London.
- MacDicken GK. 1994. Selection and management of nitrogen fixing trees. Winrock International, and Bangkok: FAO.
- Makumba W, Akinnifesi FK, Janssen B, Oenema O. 2007. Optimization of nitrogen released and immobilization from soil-applied prunings of *Sesbania sesban* and maize stover: Scientific Research and Essays. 2(9):400-407.
- Makumba WI, Akinnifesi FK, Kwesiga FR, Maghembe JA. 2002. Evaluation of *Gliricidia sepium* mixed intercropping and *Sesbania Sesban* relay cropping with maize and pigeonpea: Malawi Journal of Agricultural Sciences. 1:66-72.
- Mbuya LP et al. 1994. Useful trees and shrubs for Tanzania: Identification, Propagation and Management for Agricultural and Pastoral Communities. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- NFTA. 1994. *Sesbania sesban* : widely distributed multipurpose nitrogen Fixing Tree. NFTA 94-06. Waimanalo.
- Noad T, Birnie A. 1989. Trees of Kenya. General Printers, Nairobi.
- Owuor BO, Gudu S, Niang A. 2001. Direct seeding of *Sesbania sesban* for green manure in agroforestry systems - a short communication: Agroforestry Systems. 52(1):23-25.

Palmer E, Pitman N. 1972. *Trees of Southern Africa* Vol. 2. A.A. Balkema Cape Town.

Sileshi G, Hailu G, Mafongoya PL. 2006. Occupancy-abundance models for predicting densities of three leaf beetles damaging the multipurpose tree *Sesbania sesban* in eastern and southern Africa. Nairobi Kenya: CAB International. *Bulletin of Entomological Research*. 96:61-69.

Sileshi G, Hailu G. 2006. Modelling the spatial patterns and interspecific interactions between three chrysomelid beetles defoliating the multipurpose agroforestry tree *Sesbania sesban* in Africa: *African Entomology*. 14(2):337-348.

Sileshi G, Kenis M. 2001. Survival, longevity and fecundity of overwintering *Mesoplatys ochroptera* Stal Coleoptera: Chrysomelidae defoliating *sesbania sesban* Leguminosae and implications for its management in southern Africa: *Agricultural and Forest Entomology*. 3(3):175-181.

Sileshi G, Mafongoya PL, Kwesiga F, Kenis M, Rao MR, Maghembe JA.. 2004. The pest spectrum of *Sesbania sesban* in southern and eastern Africa: Proceedings of the Regional Agroforestry Conference on Agroforestry Impacts on Livelihoods in Southern Africa: Putting Research into Practice. p. 245-255.

Sileshi G, Mafongoya PL, Kwesiga F. 2002. Performance of *sesbania sesban* infested by the defoliating beetle *mesoplatys ochroptera* in Zambia: *Agroforestry Systems*. 56(2):177-184.

Stahl L, Hogberg P, Sellstedt A, Buresh R. 2005. Measuring nitrogen fixation by *Sesbania sesban* planted fallows using 15N tracer technique in Kenya. Umea, Sweden: Swedish University of Agricultural Science. *Agroforestry Systems*. 65:67-79.

t Mannerje L, Jones RM. 1992. *Plant Resources of South-East Asia*. No. 4: Forages. Pudoc Scientific Publishers, Wageningen.

Tietema T, Merkesdal E and Schroten J. 1992. Seed germination of indigenous trees in Botswana. Acts Press.

Vogt K. 1995. A field guide to the identification, propagation and uses of common trees and shrubs of dryland Sudan. SOS Sahel International (UK).

SUGGESTED CITATION

Orwa C, A Mutua, Kindt R , Jamnadass R, S Anthony. 2009 *Agroforestry Database: a tree reference and selection guide* version 4.0 (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)