

**LOCAL NAMES**

English (Bahamian holly, Florida holly, christmasberry tree, broadleaf pepper tree, Brazilian pepper tree); French (poivrier du Bresil, faux poivrier); German (Brasilianischer Pfefferbaum); Spanish (pimienta de Brasil, copal)

**BOTANIC DESCRIPTION**

*S. terebinthifolius* is a small tree, 3-10 m tall (occasionally up to 15 m) and 10-30 cm diameter (occasionally up to 60 cm). *S. terebinthifolius* may be multi-stemmed with arching, not drooping branches.

Leaves pinnate, up to 40 cm long, with 2-8 pairs of elliptic to lanceolate leaflets and an additional leaflet at the end. Leaflets glabrous, 1.5-7.5 cm long and 7-32 mm wide, the terminal leaflet larger than lateral ones. Leaf margins entire to serrated and glabrous.

Flowers white, in large, terminal panicles. Petals oblong to ovate, 1.2-2.5 mm long.

Fruits globose, bright red drupes, 4-5 mm in diameter.

This is a highly invasive species that has proved to be a serious weed in South Africa, Florida and Hawaii. It is also noted as invasive in other Caribbean and Indian Ocean islands. Rapid growth rate, wide environmental tolerance, prolific seed production, a high germination rate, seedling tolerance of shade, attraction of biotic dispersal agents, possible allelopathy and the ability to form dense thickets all contribute to this species' success in its exotic range.

**BIOLOGY**

*Schinus terebinthifolius* is dioecious, has high ecological plasticity, short life cycle and very rapid growth. First seed production may occur at 3 years. The flowers are insect pollinated and seed production is high. Flowering occurs in September to early November. Fruit ripening follows immediately between December and February. Seed dispersal is by animals, particularly birds and mammals including raccoons and possums which account for a major component of dispersal in the USA. Water and gravity are minor dispersal agents.



Tree; taken at: Los Angeles County Arboretum - Arcadia, CA and The National Arboretum - Washington, DC (W. Mark and J. Reimer)



Bark; taken at: Los Angeles County Arboretum - Arcadia, CA and The National Arboretum - Washington, DC (W. Mark and J. Reimer)



Fruit; taken at: Los Angeles County Arboretum - Arcadia, CA and The National Arboretum - Washington, DC (W. Mark and Reimer)

**ECOLOGY**

Brazilian pepper tree is particularly found on forest borders and river margins. It is associated with damp soils and riparian forest habitats, although it may also appear as a dry savannah plant.

**BIOPHYSICAL LIMITS**

Altitude: 0-2000 m

Temperature: 12-26°C

- Mean maximum temperature of hottest month: 20-28°C

- Mean minimum temperature of coldest month: 8-24°C

- Absolute minimum temperature: -6-3°C

Rainfall: 950-2200 mm

Soil type: prefers light, medium, heavy, well drained soils; tolerates seasonal water logging and acidity.

**DOCUMENTED SPECIES DISTRIBUTION**

Native: Argentina, Bolivia, Brazil, Chile, Paraguay, Uruguay

Exotic: Australia, Bahamas, Bermuda, Botswana, Cuba, Israel, Jamaica, Mauritius, Mexico, New Zealand, Puerto Rico, Reunion, South Africa, US, Virgin Islands (US)



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

**PRODUCTS**

**Food:** The fruits are highly appreciated as a condiment in Europe, where they are used as a substitute for black pepper (*Piper nigrum*).

**Timber:** the Brazilian pepper tree does not have significant commercial value in Brazil, but is generally used for posts, round wood; stakes; pit props; pulp and short fibre pulp.

**Fuel:** the wood is also used for fuelwood and charcoal.

**Gum or resins:** the tree is a source of resins

**Tannin or dyestuff:** also a source of tannins

**Fodder:** Brazilian pepper tree produces good quality fodder, especially for goats, but it must be used carefully because of the toxicity of some of the plant parts. The fruits are eaten by birds including American robins, and mammals, both raccoons and possums.

**Essential oils:** Essential oils extracted from the seeds of *S. terebinthifolius* have pesticidal activity against the housefly (*Musca domestica*) and its antimicrobial properties have also been investigated.

**Apiculture:** Brazilian pepper honey is highly valued especially in Florida, ensuring high incomes for beekeepers.

**Poison:** *S. terebinthifolius* may cause allergies in sensitive people even without direct contact with its leaves and fruits. For some birds, the fruit is toxic.

**Medicine:** *S. terebinthifolius* is well known for its medicinal characteristics

**SERVICES**

**Erosion control:** The tree has been used to stabilize sand dunes on the Brazilian Atlantic coast.

**Ornamental:** *S. terebinthifolius* has an attractive appearance and has therefore been introduced in many areas as an ornamental and street tree.

**Boundary or barrier:** The tree is also used in hedges.

**Reclamation:** *S. terebinthifolius* has a high ecological plasticity, a short life cycle and very rapid growth; it is therefore recommended for the restoration of degraded areas and especially gallery forests.

**TREE MANAGEMENT**

Since Brazilian pepper tree can be invasive if not properly managed, application of appropriate mechanical, chemical and biological measures should be employed to check its wanton spread.

**GERMPLASM MANAGEMENT**

Seed storage orthodox and seeds can maintain viability for about 5 months.

**PESTS AND DISEASES**

Common fungal diseases on this plant include armillaria root rot and Eutypa dieback. Pests include whitefly, citrus pyralid, citrus weevil, red-banded thrips, citrus flat mite, reniform nematode and Brazilian peppertree seed chalcid.

**FURTHER READNG**

- Austin DF. 1978. Exotic plants and their effects in southeastern Florida. *Environmental Conservation* 5(1).
- Bennett FD et al. 1988. Brazilian Peppertree -Prospects for biological control. Proc. VII Int. Symp. Biol. Contr. Weeds, 6-11, March 1988, Rome, Italy.
- Binggeli P. 1998. An Overview of Invasive Woody Plants in the Tropics. School of Agricultural and Forest Sciences Publication Number 13, Unversity of Wales, Bangor.
- Buss CM. 2002. The potential threat of invasive tree species in Botswana. Department of Crop Production and Forestry, Ministry of Agriculture, Government of Botswana.
- Carvalho PER. 1994. Brazilian forest tree species: silvicultural recommendations, potential and use of wood. Brasília, Brazil.
- Cassani JR. 1986 Arthropods on Brazilian Peppertree, *Schinus terebinthifolius* (Anacardiaceae), in south Florida. *Florida Entomologist* 69(11).
- Cronk QCB & Fuller JL. 1995. Plant invaders: the threat to natural ecosystems. Chapman & Hall, London, UK.
- Ferriter A & Clark D. 1997. Brazilian pepper management plan for Florida. Brazilian Pepper Task Force, Florida Exotic Pest Plant Council.
- Gogue GJ, Hurst CJ & Bancroft L. 1974. Growth inhibition by *Schinus terebinthifolius*. *HortScience* 9(3).
- Hasui E & Hofling E. 1998. Food preference of the frugivorous birds of semideciduous forest, Sao Paulo, Brazil. *Iheringia - Série Zoologia*.
- Henderson L. 2001. Alien Weeds and Invasive Plants. Plant Protection Research Institute Handbook No. 12. Paarl Printers, Cape Town, South Africa.
- Kunkel G. 1978. Flowering trees in subtropical gardens. Dr. W. Junk b. v. Publishers, The Hague.
- Laca-Buendía JP, Brandao M, Oliveira LMda S. 1992. Utilization of the fruits of *Schinus terebinthifolius* Raddi. (Anacardiaceae) as a substitute for black pepper (*Piper nigrum* L.). *Daphne, Revista do Herbário PAMG da EPAMIG*.
- Little ELJr, Woodbury RO, Wadsworth FH. 1974. Trees of Puerto Rico and the Virgin Islands. Second volume. Agriculture Handbook, US Department of Agriculture, No. 449.
- Luken JO & Thieret JW. 1997. Assessment and Management of Plant Invasions. Springer-Verlag, New York, USA.
- Martínez MJ, Betancourt J, Alonso-González N, Jauregui A. 1996. Screening of some Cuban medicinal plants for antimicrobial activity. *Journal of Ethnopharmacology*, 52(3).
- Morton JF. 1978. Brazilian pepper (*Schinus terebinthifolius*): its impact on people, animals and the environment. *Economic Botany*.
- Olmsted I & Yates S. 1984. Florida's pepper problem. Garden, USA
- Owen SJ. 1996. Ecological weeds on conservation land in New Zealand: A database. Department of Conservation, DOC Science Publications, Wellington, New Zealand.
- Phillips S. 1981. Memo: Brazilian Pepper control master plan and research project. The Sanibel Captiva Conservation Foundation, Inc. Sanibel, FL.
- Pirani JR & Cortopassi-Laurino M. 1993. Flores e Abelhas em Sao Paulo. Sao Paulo, Brazil.
- Randall J. 2003. *Schinus terebinthifolius*. California Invasive Pest Plant Council (Cal-IPPC), USA.
- Siddiqui RR, Ahmad H, Shakoor ChS, Ehteshamuddin AFM, Shireen S. 1996. Antimicrobial activity of essential oils. Part II. *Pakistan Journal of Scientific and Industrial Research*, 39(1/4).
- Tomlinson PB. 1980. The biology of trees native to tropical Florida. Printed and bound by Harvard University Printing Office, Allston, Mass.
- Uphof JC Th. 1968. Dictionary of Economic Plants. Stechert-Hafner Service Agency Inc., New York, New York.
- Weber E. 2003. Invasive Plant Species of the World. A Reference Guide to Environmental Weeds. CABI Publishing, Wallingford, UK.
- Westbrooks RG. 1998. Invasive plants, changing the landscape of America: Fact book. Washington DC, USA: Federal Interagency Committee for the Management of Noxious and Exotic Weeds (FICMNEW).
- Woodall SL. 1982. Herbicide tests for control of Brazilian-Pepper and *Melaleuca* in Florida. *Southeastern Forest*

Experiment Station, Asheville, N. C. USDA Forest Service Research Note.

Wunderlin RP. 1982. Guide to the vascular plants of central Florida. Univ. Presses Gainesville, Florida.

**SUGGESTED CITATION**

Orwa C, Mutua A , Kindt R , Jamnadass R, Simons A. 2009. Agroforestry Database:a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/af/treedb/>)