

## Cupressus lusitanica

cypress

### LOCAL NAMES

Amharic (yeferenji-tid); English (cypress, cedar of Goa, Mexican cypress, Kenya cypress); French (cyprès, cyprès de Mexico, cyprès de Goa); German (Mexikanische Zypresse, blaugrüne Zypresse); Spanish (ciprés Mexicano, ciprés de Portugal, ciprés); Swahili (msanduku); Tigrigna (tsehdiferenji); Trade name (cypress)

### BOTANIC DESCRIPTION

*Cupressus lusitanica* is an evergreen tree, 35 m high, with a dense, conical crown. Branches spread out widely but terminate in pendulous branchlets. Trunk short, 70 cm in diameter. Bark on trunk is reddish-brown, exfoliating in long, narrow strips, eventually becoming roughened by the development of many short cracks. Branchlets covered with small, decussate, adpressed, acute, sessile, scale leaves with flexed tips.

The distinctly bluish-green foliage is ovate, closely pressed, usually with long, pointed apex.

Male flowers small, oblong or cylindrical; female subglobose, very small, composed of 6-14 fertile decussate scales with several ovules each.

Cones ellipsoid; bluish-green when young turning reddish-brown when mature; 13-25 mm in diameter; composed of 6-12 woody, shieldlike scales, each bearing 8-10 seeds; male cones appear to be fat tips to branchlets and produce clouds of yellow pollen; female cones rounded, scales with central pointed projections. Seeds brown, with resin glands, up to 4 mm long, with a narrow wing.

The Latin name 'Cupressus' comes from the Greek 'kuparessos', which commemorates a youth of that name who was turned into a cypress tree by Apollo. The specific name is derived from Lusitania, Portugal, where the tree was introduced in the 17th century.

### BIOLOGY

Trees 1st bear fruit at 6-9 years, or later on unfavourable sites. Flowering takes place at the driest time of the year, with male and female flowers arising at different points on the crown (monosexual). Cones develop within 6 months after wind pollination and take 2 years to mature.

Mill.

Cupressaceae



*C. lusitanica* windbreaks in dairy pasture, Volcan Barba, Costa Rica. (David Boshier)



*C. lusitanica* unthinned plantation, Cartago, Costa Rica. (David Boshier)



*C. lusitanica* young seedlings in nursery. (David Boshier)

# Cupressus lusitanica

Mill.

Cupressaceae

cypress

## ECOLOGY

*C. lusitanica* is found in seasonally moist to permanently moist climates, with annual precipitation typically between 1000 and 1500 mm and a dry season lasting not more than 2-3 months. It also occurs in very moist climates with annual precipitation up to 4000 mm. It is not generally damaged by occasional snow or brief periods of frost, but there are significant differences in this among provenances. It cannot withstand waterlogging. Associated species in its natural habitat include *Abies guatemalensis*, *Litsea glaucescens*, *Pinus montezumae* and *Prunus brachybotrya*.

## BIOPHYSICAL LIMITS


Altitude: 1 000-4 000 m, Mean annual temperature: 12-30 deg. C, Mean annual rainfall: 800-1 500 mm

Soil type: *C. lusitanica* flourishes in deep, moist, well-drained, fertile loams of neutral to slightly acidic reaction.

## DOCUMENTED SPECIES DISTRIBUTION

Native: Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama, US

Exotic: Eritrea, Ethiopia, Kenya, Portugal, South Africa, Spain, Tanzania, Uganda



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

## **Cupressus lusitanica**

cypress

Mill.

Cupressaceae

---

### **PRODUCTS**

Fuel: *C. lusitanica* is a good source of firewood.

Timber: The white wood saws cleanly and has straight fine grain; it is a source of construction wood and pulp wood and is used for furniture, poles and posts.

### **SERVICES**

Shade or shelter: Trees are suitable as windbreaks.

Ornamental: The beautiful tree can be planted in amenity areas.

Boundary or barrier or support: It is grown as a live fence.

**TREE MANAGEMENT**

The fast-growing conifers are planted on land that has been cleared by burning, to improve rooting conditions and eliminate potential competitors. In general, the seedlings should be planted at 2 x 2 or 3 x 3 m; close spacing is preferable to prevent strong branches from developing. As *C. lusitanica* gives only limited protection against soil erosion, pure stands on slopes or erosion-prone sites should be underplanted with other suitable species. Weeding is an absolute must during the 1st years.

Pruning is practised so that trees yield high-quality, knot-free sawlogs. Pruning should be done when trees are 3 years old, with 3 further prunings at 6, 9 and 13 years of age in a 25-30 year rotation. Trees grown for high-quality timber should be pruned to 30% of their stem height every 3 years without diminishing volume growth. Thinning is prescribed before each pruning.

In early years, individual trees should show an annual height increment of 1.2-1.5 m (2 m in exceptional cases). Trees produce poles after 10 years and general-purpose timber after 20 years. They need to be protected from fire and rodent attack.

**GERMPLASM MANAGEMENT**

After extraction, the seeds are dried in the sun to 6-10% mc before they are stored. Long-term seed storage behaviour is orthodox. Germination rate for seeds in hermitic storage for 21 years at 5 deg. C with less than 10% mc is 10%; viability can be maintained in hermitic storage at 3 deg. C with 6-10% mc. With normal dry storage, 75% germination can be maintained for 1 year and for several years if seeds are refrigerated at 4 deg. C. Seed weight varies with provenance, ranging from 170 000 to 320 000 seeds/kg.

**PESTS AND DISEASES**

Trees are susceptible to insect pests *Acheta assimilis*, *Agrotis* spp., *Atta* spp., *Captotermes crassus*, *Exophthalmus* spp., *Phytophaga* spp., *Platypus* spp. and *Pypselonotus aratus*. The pest most widely known is the timber borer *Oemida gahani*, which gains entry through surface wounds such as those created by pruning and degrades timber; pruning before trees are 7 years old could contain the problem. The cypress aphid *Cinara cupressi* attacks trees of the Cupressaceae family in south and central Africa and is fast advancing to Tanzania and Uganda. Pests are controlled using insecticides and biological control techniques, by planting resistant varieties and by legislation prohibiting entry of plant material from infested areas.

The most serious diseases are those affecting cambial function and other tissues. In misty and wet weather, trees are susceptible to fungal attack by *Monochoetia unicornis*. It infects stems of young plants and young shoots of older trees and appears as cankers and local disorganization of newly formed bark, cambial and cortical tissues. *Armillaria* root rot caused by *Armillaria mellea* attacks roots and butts of trees and causes rotting, which eventually kills the trees. Pathogens that attack trees include *Cercospora sequoiae*, *Colletotrichum* spp., *Fusarium* spp., *Seridium cardinale* and *Verticillium*.

**FURTHER READING**

- Albrecht J. ed. 1993. Tree seed hand book of Kenya. GTZ Forestry Seed Center Muguga, Nairobi, Kenya.
- Asfaw Z. 1969. Silvicultural management of *Cupressus lusitanica* and *Grevillea robusta* in southern Ethiopia. M. Phil. Research Outline and Progress Report. School of Agricultural and Forest Sciences U.C.N.W., Bangor.
- Bein E. 1996. Useful trees and shrubs in Eritrea. Regional Soil Conservation Unit (RSCU), Nairobi, Kenya.
- Bekele-Tesemma A, Birnie A, Tengnas B. 1993. Useful trees and shrubs for Ethiopia. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Birnie A. 1997. What tree is that? A beginner's guide to 40 trees in Kenya. Jacaranda designs Ltd.
- Hong TD, Linington S, Ellis RH. 1996. Seed storage behaviour: a compendium. Handbooks for Genebanks: No. 4. IPGRI.
- ICRAF. 1992. A selection of useful trees and shrubs for Kenya: Notes on their identification, propagation and management for use by farming and pastoral communities. ICRAF.
- Katende AB et al. 1995. Useful trees and shrubs for Uganda. Identification, Propagation and Management for Agricultural and Pastoral Communities. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Mbuya LP et al. 1994. Useful trees and shrubs for Tanzania: Identification, Propagation and Management for Agricultural and Pastoral Communities. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Noad T, Birnie A. 1989. Trees of Kenya. General Printers, Nairobi.
- Odera J. 1990. Know and prevent entry of the cypress aphid (*Cinara cupressi*) into Kenya. KEFRI Technical No. 7.

**SUGGESTED CITATION**

Orwa C, Mutua A, Kindt R, Jamnadass R, Simons A. 2009. Agroforestry Database: a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/af/treedb/>)