Equador,	Peru
----------	------

**Common names:** Eng: cherimola, cherimoya, custard apple; Swa: mtomokc mtopetope.

Ecology: A small tree growing at 800 m or higher in the Andes. The well-flavoured refreshing fruit was known to the ancient people of the area. It cannot tolerate a hot lowland climate and is grown commercially in Chile, Spain, USA and New Zealand. In East Africa the name "custard apple" has been used loosely for Annona species and hybrids. They grow best in cooler tropical uplands over 1,000 m, as in Lushoto, in sandy loam soils with a dry season. They tolerate lower temperatures for growth and fruiting than *A. squamosa*. Food (fruit).

A deciduous tree 3-10 m, often branched from the base. **Description:** LEAVES: Alternate, light green, oval and pointed, sometimes narrow, 12-20 cm long withiew hairs above but characteristic golden-brown hairs below. (Young leaves and veins on mature leaves.) FLOWERS: fragrant, usually solitary (or 2-3) hang down on short hairy stalks, 3 outside petals narrow to 3 cm, pale vellow with a purple spot at the base, 3 inner petals red-purple, tiny. FRUIT: Green and compound, 8-15 cm across, variable, heart-shaped, round or oval with regular overlapping fleshy scales or the surface patterned with U-shaped depressions, spirally arranged like fingerprints. The ripe soft flesh is creamwhite, granular, acid-sweet, easily separated from numerous seeds, each one brown, about 1.5 cm long. All Annonas can be grown from seed and cuttings, but **Propagation:** budding and grafting are more successful-with A. cherimoya or A. squamosa as rootstock. Seed info.: Similar to A. squamosa.

Seed info.: treatment: storage:

Management:Small beetles usually pollinate the flowers. Hand pollination<br/>improves fruit set and the fruit ripens 4-5 months later.Remarks:Green fruit, seed and leaves of Annona spp. are reported to<br/>have insecticidal properties. Fruit and seed are effective<br/>against worms. Fruit are eaten fresh and the pulp can be<br/>used to flavour ice cream. The hybrids between A. cherimola<br/>and A. squamosa are known as atemoya or custard apple<br/>and are grown all over Africa and the sub-tropics. The<br/>word "annona" means yearly production and indicates a<br/>good annual yield of fruit.

