Callistemon citrinus var. splendens

Myrtaceae

Australia, New Zealand, Tasmania

| Common names: Ecology: | Eng: bottlebrush tree, weeping bottlebrush. In Australia it is found on low-lying ground including valley bottoms, so will grow on poorly drained sites. Now widely planted in the tropics as an ornamental. |
|---------------------------|---|
| Uses: | Firewood, charcoal, bee forage, windbreak, ornamental. |
| Description: | A small evergreen tree, to 6 m high, with drooping foliage. BARK: grey, smooth, furrowed with age. LEAVES: narrow, tough, grey-green to 8 cm, young leaves pink- green, faintly lemon scented when crushed. FLOWERS: vivid crimson bottlebrush-like cylindrical spikes, a mass of long red stamens, nectar attracting sunbirds and bees. Leafy shoots continue to grow beyond the flower "brush". FRUIT: small woody capsules, persisting many months, contain the tiny seed. |
| Propagation: | Seedlings. |
| Seed info.: | No. of seeds per kg: about 44,000. Seed germination is uniform (two weeks). |
| treatment: storage: | not necessary. seed stores well (six months). |
| Management: | Fast growing on good sites. |
| Remarks: | Commonly planted in gardens. Another Callistemon variety has erect flowers, not drooping, otherwise it is similar. |

Callistemon citrinus var. splendens

Myrtaceae

